

# WD60/FAP163 is a Dynein Intermediate Chain Required for Retrograde Intraflagellar Transport in Cilia

Ramila S. Patel-King<sup>+\$</sup>, Renée M. Gilberti<sup>+\$†</sup>, Erik F.Y. Hom<sup>#</sup>  
and  
Stephen M. King<sup>+\*</sup>

<sup>+</sup> Department of Molecular, Microbial and Structural Biology,  
University of Connecticut Health Center  
263 Farmington Avenue  
Farmington, Connecticut 06030-3305

and

<sup>#</sup> Department of Molecular and Cellular Biology,  
Harvard University,  
520 Oxford Street, NW 469  
Cambridge, Massachusetts 02138

Tel: (860) 679 3347

Fax: (860) 679 3408

Email: [king@neuron.uhc.edu](mailto:king@neuron.uhc.edu)

\* To whom correspondence should be addressed

<sup>\$</sup> R.S.P-K and R.M.G. contributed equally

<sup>†</sup> Present address: Center for Academic Programs, University of Connecticut, 368 Fairfield Way,  
Unit 4170, Storrs, CT 06269-4170.

Key Words: Cilia / Dynein / Flagella / Intraflagellar Transport / Microtubules

Running Title: WD60/FAP163 is required for Ciliary Assembly

## Abstract

Retrograde intraflagellar transport (IFT) is required for assembly of cilia. We identified a *Chlamydomonas* flagellar protein (FAP163) as being closely related to the D1bIC(FAP133) intermediate chain of the dynein that powers this movement. Biochemical analysis revealed that FAP163 is present in the flagellar matrix and is actively trafficked by IFT. Furthermore, FAP163 co-purified with D1bIC(FAP133) and the LC8 dynein light chain indicating that it is an integral component of the retrograde IFT dynein. To assess the functional role of FAP163, we

generated an RNAi knockdown of the orthologous protein (WD60) in planaria. The *Smed-wd60(RNAi)* animals had a severe ciliary assembly defect that dramatically compromised whole organism motility. Most cilia were present as short stubs that had accumulated large quantities of IFT particle-like material between the doublet microtubules and the membrane. The few remaining approximately full-length cilia had a chaotic beat with a frequency reduced from 24 to ~10 Hz. Thus, WD60/FAP163 is a dynein intermediate chain that is absolutely required for retrograde IFT and ciliary assembly.

### **Introduction**

Intraflagellar transport (IFT <sup>footnote 1</sup>) is a complex bidirectional motility system that is required for the assembly and maintenance of both motile and primary/sensory cilia (Rosenbaum and Witman, 2002). In general, the components of this system have been broadly conserved from protists to vertebrates and defects impact a wide array of motility, signaling and sensory functions; in vertebrates, this leads to many different developmental defects and complex syndromes *e.g.* (Pazour *et al.*, 2000; Li *et al.*, 2004; Hildebrandt *et al.*, 2011).

The core of the IFT system consists of particles (or trains) that are comprised of multiple copies of two subcomplexes (termed A and B) that in turn have very distinct protein compositions (Cole, 2003). A combination of genetics, biochemistry and fluorescent imaging has suggested that IFT-B is necessary for anterograde transport whereas IFT-A functions in the retrograde direction (reviewed in (Cole, 2009)). Cargoes required for assembly and maintenance of cilia are thought to be loaded onto the IFT trains at the basal body region of the cell. Following gated entry into the ciliary compartment, these IFT particles are then transported towards the ciliary tip by one or more kinesin motors (Snow *et al.*, 2004). Following reorganization at the tip, IFT particles are returned to the ciliary base by a dynein that is more closely related to canonical cytoplasmic dynein than it is to those that power the axoneme.

Mutants defective in known components of this retrograde motor are able only to assemble short ciliary stubs in which the region between the doublet microtubules and the membrane is swollen due to the accumulation of IFT particles that are sent into the organelle but cannot be returned to the cell body (Pazour et al., 1999; Signor et al., 1999).

Two different kinesins have been implicated in anterograde IFT in several organisms *e.g.* (Wickstead and Gull, 2006; Pan *et al.*, 2006 ), however it has been generally assumed that there is a single dynein holo-enzyme responsible for retrograde transport (here termed the IFT dynein for clarity though it may have additional functions outside of cilia). However, there has been a suggestion that two different dyneins might participate in this process in *Caenorhabditis elegans* (Hao et al., 2011). In *Chlamydomonas*, the IFT dynein is thought to contain two HC motor units (D1bHC encoded at *DHC16*) (Pazour et al., 1999; Porter et al., 1999) that are tightly associated with light intermediate chains (D1bLIC encoded at *DLII*) (Perrone et al., 2003) and an IC/LC complex that consists of a WD-repeat intermediate chain [D1bIC(FAP133) encoded at *DIC5*] (Rompolas et al., 2007) and the LC8 dynein light chain dimer (encoded at *DLL1* a.k.a. *FLA14*) (Pazour et al., 1998). Canonical cytoplasmic dynein also contains LC dimers of the DYNLT (Tctex1) and DYNLRB (roadblock) families; although these have not yet been identified biochemically within the *Chlamydomonas* IFT dynein (Witman, 2012), there is a evidence that a DYNLT ortholog may be part of this complex in *Ca. elegans* (Efimenko et al., 2005). Detailed mechanistic analysis of the IFT dynein has proven rather difficult as it tends to readily dissociate into at least two different subparticles during purification (Rompolas et al., 2007). One consists of the HC motor and D1bLIC (Pazour *et al.*, 1999; Porter *et al.*, 1999; Perrone *et al.*, 2003), whereas the second contains D1bIC(FAP133) and LC8 (Pazour et al., 1998; Rompolas et al.,

2007). However, it has remained unclear whether this composition represents the complete particle or if additional proteins or indeed dynein subtypes are required for retrograde transport.

Here we identify an additional conserved WD-repeat protein that is closely related to D1bIC(FAP133) (the orthologous protein is termed WD34/WDR34 in mammals). This protein (FAP163)<sup>footnote2</sup> is present in the *Chlamydomonas* flagella proteome (Pazour et al., 2005) and its orthologue WD60/WDR60 was upregulated in microarray data for ciliated epithelia in mouse trachea (Hoh et al., 2012), suggesting that it may be involved in IFT. We demonstrate here that *Chlamydomonas* FAP163 is present in the flagellar matrix, associates with D1bIC(FAP133) and is actively trafficked by IFT. Furthermore, we have obtained an RNAi-mediated knockdown of WD60 expression in the planarian flatworm *Schmidtea mediterranea* and find that this protein is indeed absolutely required for ciliary assembly and that its lack results in the failure of retrograde IFT.

## Results

### Bioinformatics of Dynein-associated WD-repeat Proteins

Alignment of dynein IC and related WD-repeat protein sequences by the Panther classification algorithm (PTHR12442) identified a flagellar-associated *Chlamydomonas* WD-repeat protein of unknown function [FAP163; (Pazour *et al.*, 2005)] as being most closely related to D1bIC(FAP133) that we previously demonstrated was an IC of the dynein complex responsible for retrograde IFT in *Chlamydomonas* (Rompolas *et al.*, 2007). Further analysis using BLAST searches and multiple sequence alignments revealed orthologs of FAP163 (termed WD60/WDR60 in *H. sapiens*) in a broad range of organisms that assemble cilia including mammals and other vertebrates, sea urchins, various protists and the planarian *Schmidtea mediterranea* (mk4.000489.09.01; Ψ-BLAST score with human WD60 after two iterations =

$9 \times 10^{-103}$ ) (Fig. 1a). Sequence analysis of human WD60 (1066 residues; Mol. Wt. = 122,570 Da; pI = 6.88) revealed that it consists of an N-terminal segment that is predicted to be almost completely  $\alpha$ -helical, containing two short stretches that have a very high probability of forming coiled coils (Fig. 1b) and a repetitive region (residues 24-328) containing many charged residues (Fig. 1c), followed by a C-terminal region that is predicted to consist entirely of a series of  $\beta$  strands with a single short helical segment near the very terminus. Three canonical WD repeats were found within this  $\beta$ -strand segment using the SMART algorithm (Fig. 1d); at least three additional putative degenerate WD repeats were tentatively identified by visual inspection of the predicted secondary structure; one is located prior to the first canonical WD repeat and two more the first and second canonical repeats. In addition, a single IQ motif (residues 432-450) was identified by the Eukaryotic Linear Motif Resource, suggesting that this protein may bind calmodulin.

### **Biochemical Analysis of FAP163 in *Chlamydomonas* Flagella**

Analysis of the *Chlamydomonas* flagellar proteome revealed the presence of FAP163 migrating at  $M_r \sim 90,000$  (Pazour *et al.*, 2005), although the calculated mass is only 76,134 Da. When blot-purified antibody CT295 raised against the FAP163 C-terminal domain was used to probe samples of wildtype flagella, a single major band of  $M_r 89,000$  was detected that is completely consistent with the electrophoretic migration of FAP163 observed in the proteomic study (Fig. 2a). FAP163 was present in flagella from mutants that lack outer arms (*oda1-6*), inner arms (*ida1*, *ida4*), radial spokes (*pf14*) or the central pair microtubule complex (*pf18*) (Fig. 2b), although levels did appear reduced in *ida1* and possibly *ida4*. To determine where FAP163 is located within the flagellum, membrane/matrix components were solubilized by detergent treatment; FAP163 was observed almost exclusively in the soluble fraction. Similarly, most

FAP163 was also present in the extract obtained from flagella following multiple rounds of freeze/thaw which releases the flagellar matrix but leaves the membrane associated with the axoneme (Fig. 2c). Thus, FAP163 is a component of the flagellar matrix and is not tightly bound to the microtubular axoneme or the membrane.

To assess the oligomeric state of FAP163, freeze-thaw extracts were fractionated by chromatography in a Superose 6 gel filtration column. FAP163 was found to elute in a single broad peak with an estimated mass of >350 kDa, and precisely co-purified with one peak of the LC8 dynein light chain that is present in multiple flagellar complexes (Fig 3a). Furthermore, we observed that under these solution conditions, FAP163 also co-eluted with D1bIC(FAP133) suggesting that both WD-repeat proteins might be part of the same complex. To further test this putative association, a wildtype freeze/thaw flagella extract was fractionated by ion exchange chromatography in a MonoQ column (Fig. 3b). D1bIC(FAP133) was found in two distinct peaks as described previously (Rompolas et al., 2007). We also observed that FAP163 precisely co-purified with both D1bIC(FAP133) peaks. These data strongly suggest that FAP163 and D1bIC(FAP133) are associated within the same IC/LC complex

To determine whether FAP163 is transported by the IFT machinery, wildtype (cc125) and *fla10* cells were incubated at 22 or 32°C for three hours and then deflagellated. As expected, analysis of the resulting flagella revealed that D1bIC(FAP133), a component of the retrograde IFT motor, was highly depleted in flagella from *fla10* cells held at the restrictive temperature compared to wildtype. Similarly, we observed that FAP163 levels also were reduced whereas those of the axoneme-associated outer arm dynein proteins LC1 and IC1 were not (Fig. 4). This result indicates that flagellar FAP163 is actively trafficked by IFT.

### **RNAi-mediated Knockdown of WD60 in Planaria**

No *Chlamydomonas* FAP163 mutant currently exists. Therefore, to test the hypothesis that WD60/FAP163 might represent another dynein IC that plays a key role in retrograde IFT, we performed an RNAi-mediated knockdown in the planarian *S. mediterranea*; interestingly, this organism lacks a recognizable D1bIC(FAP133). Accordingly, we obtained a 325 bp region of the WD60 coding region by PCR from planarian first-strand cDNA and subcloned this into the L4440 vector that has opposing T7 promoters, and expressed dsRNA in *E. coli* strain HT115(DE3). Planaria were fed every 2-3 days for several weeks with this dsRNA or with bacteria containing an empty vector.

Analysis of mRNA by RT-PCR, from first-strand cDNA prepared using a random hexamer as the primer, revealed that two bands for WD60 were present. One product obtained was the originally targeted 325 bp fragment that encodes the central portion of the WD60 protein, whereas the second product of 502 bp consisted of this same exonic segment but included two unspliced introns (Fig. 5a, b). The 502 bp product was not observed when first-strand cDNA was prepared using oligo-dT as the primer for reverse transcriptase, nor was it obtained when the original RNA sample was treated with DNaseI. Consequently, this second product derives from genomic DNA. Interestingly, when first-strand cDNA was prepared using an oligo-dT primer, we obtained the 325 bp product and two minor additional products that included one or other of the two introns (fig. 5c). Thus, planaria maintain a detectable pool of partially-spliced polyadenylated mRNA for WD60. RT-PCR using total RNA derived from *Smed-wd60(RNAi)* planaria further revealed that ingestion of WD60 dsRNA resulted in greatly decreased amounts of the 325 bp product compared to empty vector or no vector controls, but had essentially no effect on transcript levels for actin, IFT88 or IC2 (Fig. 5a).

### **Phenotypic Analysis of *Smed-wd60(RNAi)* Planaria**

We next monitored organismal gliding locomotion and found that *Smed-wd60(RNAi)* animals moved at  $0.55 \pm 0.19$  mm/s (mean  $\pm$  standard deviation; n = 13) compared to  $1.48 \pm 0.21$  mm/s (n = 7) for empty vector controls (Fig. 5d, e and movie S1). This decrease in gliding velocity is similar to that observed for planaria which lack functional outer dynein arms (*Smed-ic2(RNAi)*) or indeed are unable to assemble cilia at all (*Smed-ift88(RNAi)*) (Rompolas *et al.*, 2010). Furthermore, *Smed-wd60(RNAi)* animals exhibited the alteration in the mode of motility, from cilia-driven gliding to peristaltic waves of muscular contraction, that was found previously (Rompolas *et al.*, 2010) to be a defining characteristic of defects in ciliary assembly and motility (Fig. 5f and movie S2). Following prolonged feeding with dsRNA for over 4 weeks, *Smed-wd60(RNAi)* planaria became essentially stationary, and had no discernable cilia at the dorsal/ventral margin when examined by differential interference contrast microscopy. Furthermore, these planaria were unable to use the musculature in the rear of the body and eventually split approximately half way along their length leading to the partial disintegration of the animal.

Examination of the ventral surface by scanning EM of animals fed the RNAi vector for ~3 weeks revealed a dramatic reduction in both the number and length of cilia assembled by *Smed-wd60(RNAi)* planaria compared to controls (Fig. 6). In control animals, the ventral epithelium bore thousands of ~10  $\mu$ m-long cilia especially towards the anterior and in the central region surrounding the pharyngeal pore (Fig. 6a and c). In contrast, only relatively few almost full-length cilia and many short ciliary stubs (<0.5  $\mu$ m in length) were evident on *Smed-wd60(RNAi)* planaria (Fig. 6b, d and e). These observations suggested that WD60 is required for ciliary assembly. To further assess this hypothesis, we examined cilia and ciliary stubs from *Smed-wd60(RNAi)* planaria by thin-section transmission EM (Fig. 7). We found that the ciliary

stubs contained doublet microtubules and associated structures, but had an enlarged diameter due to the presence of large quantities of amorphous electron-dense material located between the axonemal doublet microtubules and the ciliary membrane. Indeed, even in longer cilia, accumulations of this material were readily evident.

High-speed video microscopy and kymograph analysis revealed that the few longer cilia remaining on the ventral surface of *Smed-wd60(RNAi)* planaria were motile (Fig. 8a, and movie S3). However, their beat appeared uncoordinated and occurred at rates of ~10-14 Hz; some cilia were observed to protrude essentially vertically from the epithelium and to move only at the base leading to a twitching motion. In contrast, the cilia of control animals exhibited a high degree of metachronal synchrony and beat at ~24 Hz with both power and recovery strokes (Fig. 8b, and movie S4).

## Discussion

In this report, we have examined the functional role of WD60/FAP163, a WD-repeat protein that has been broadly conserved amongst ciliated organisms. Sequence analysis suggested that this protein is an IC of the dynein that powers retrograde intraflagellar transport and we found that it is indeed present in the *Chlamydomonas* flagellar matrix and actively trafficked by the IFT machinery. Disruption of WD60/FAP163 function in planaria by RNAi resulted in the failure of ciliary assembly and the accumulation of IFT particle-like material in the resulting ciliary stubs with the consequent disruption of organismal motility. Previously, an essentially identical ciliary assembly phenotype in *Chlamydomonas* was demonstrated to be caused by the failure of retrograde IFT dynein function *e.g.* (Pazour *et al.*, 1999; Porter *et al.*, 1999). Thus, it appears that WD60/FAP163 is an intergral dynein IC that in planaria is absolutely required for ciliary assembly.

## **FAP163/WD60 is a Component of the IFT Machinery and Required for Retrograde IFT**

Phylogenetic analysis revealed that WD60/FAP163 groups with the dynein IC subfamily of WD-repeat proteins and indeed these proteins are most closely related to D1bIC(FAP133 or WD34) that we previously demonstrated to be part of the IFT dynein in *Chlamydomonas* (Rompolas et al., 2007). Mammalian WD60 has an N-terminal region that is highly repetitive and contains two segments of coiled coil followed by a C-terminal domain consisting of WD-repeat motifs. Furthermore, there is a single IQ motif located within the N-terminal region. Although this motif does not appear to be present in *Chlamydomonas* FAP163, it does raise the possibility that WD60 in vertebrates binds calmodulin and thus may be responsive to alterations in  $\text{Ca}^{2+}$  levels. In contrast, D1bIC(FAP133) consists of a short N-terminal domain containing two degenerate LC8 binding sites followed by a C-terminal segment consisting of six WD-repeats (Rompolas et al., 2007).

Biochemical analysis revealed that FAP163 is almost exclusively present in the flagellar matrix as are core IFT components and associated kinesin and dynein motors. Using the *fla10* temperature-sensitive anterograde IFT mutant, we found that FAP163 was reduced in *Chlamydomonas* flagella when cells were held at the restrictive temperature suggesting that this protein is indeed trafficked by IFT. In contrast, the flagellar levels of FAP163 were not affected by the presence or absence of outer arms, radial spokes or the central pair complex, although they did appear decreased in the *ida1* and perhaps also *ida4* inner arm dynein mutants. When expression of WD60/FAP163 was disrupted in planaria, we observed severe disruption of ciliary structures such that, in general, only short ciliary stubs were retained. These stubs contained accumulated material located between the outer doublet microtubules and the membrane as is commonly seen in *Chlamydomonas* strains unable to undergo retrograde IFT (Pazour *et al.*,

1998; Pazour *et al.*, 1999; Porter *et al.*, 1999). Thus, WD60/FAP163 is absolutely required for retrograde IFT. This failure of ciliary assembly on the ventral epithelium led to a change in planarian whole organism locomotion, from cilia-driven gliding to muscle-based peristaltic waves, which we demonstrated previously is characteristic of defects in ciliary assembly or motile behavior (Rompolas *et al.*, 2010).

### **Organization of the Dynein Motor that Powers Retrograde Intraflagellar Transport**

Dyneins with multiple HC motor units (*i.e.* outer arm, inner arm I1/f and canonical cytoplasmic dynein) all contain an IC/LC complex that at a minimum consists of two WD-repeat ICs associated with dimers of three distinct LC species and is absolutely required for assembly of the dynein particle *e.g.* (Tang *et al.*, 1982; Mitchell and Kang, 1991); the ICs within these complexes associate directly (Mitchell and Rosenbaum, 1986; King *et al.*, 1991; Lo *et al.*, 2006), whereas LC binding apparently brings structure to inherently disordered regions of the ICs (Nyarko *et al.*, 2004). In axonemal dyneins, the two ICs in the complex are different gene products and form obligate heterodimers (reviewed in (King, 2012)), whereas in canonical mammalian cytoplasmic dynein the various alternative spliced forms of the two very closely related IC genes are capable of forming both homo- and hetero-dimers (Lo *et al.*, 2006), although whether all the possible combinations actually occur *in vivo* remains uncertain.

Previously, the dynein that powers retrograde IFT in *Chlamydomonas* was known to contain a distinct HC and associated LIC, as well as D1bIC (also known as FAP133 or WD34) with bound LC8 dimer. Our biochemical analysis using two different chromatographic procedures to fractionate *Chlamydomonas* flagellar matrix components demonstrated that FAP163 is also part of this complex and thus FAP163 and D1bIC(FAP133) presumably form a heterodimer that is further stabilized by LC8. The similar reductions in flagellar levels of both

FAP163 and D1bIC(FAP133) in the *fla10* mutant shifted to the restrictive temperature further supports this suggestion. We also observed that FAP163 and D1bIC(FAP133) elute together in two distinct complexes following anion exchange, and we demonstrated previously that both these peaks of D1bIC(FAP133) subsequently migrate at ~7 S in sucrose density gradients (Rompolas et al., 2007) and are thus not associated with the D1bHC. This raises the possibility that there are two forms of the IC/LC complex. Further biochemical analysis will be required to determine whether these two IC/LC complex forms differ in protein content or if the change in elution properties is the result of post-translational modifications.

Intriguingly, although platyhelminths such as *Ca. elegans* and *S. mediterranea* express most components of the IFT dynein and exhibit retrograde IFT (*e.g.* (Signor et al., 1999)), they do not encode a recognizable ortholog of D1bIC (Wickstead and Gull, 2012). Given that ICs are thought to be essential for the assembly and function of dyneins with multiple motor domains, this observation was rather puzzling as it implied that the dynein mediating retrograde IFT in platyhelminths might have an organization distinct from that of all other multi-motor dyneins. Our current results suggest an alternative; namely that some organisms (such as *Chlamydomonas* and vertebrates) actually employ two distinct dynein ICs for retrograde IFT, whereas planarians appear to have secondarily lost the ortholog of D1bIC(/FAP133/WD34) and only utilize WD60/FAP163. Thus, in *Schmidtea* the IC/LC complex of the IFT dynein may be composed of a FAP163/WD60 homodimer; alternatively, it is possible that an additional IC awaits identification in this organism. Surprisingly, *Ca. elegans* also appears to lack a WD60/FAP163 ortholog; thus, there may be even more diversity in IFT dynein components. Indeed, it has been reported that a second IFT dynein based on a different HC may exist in *Ca. elegans* (Hao et al., 2011), further suggesting that the mechanisms of retrograde IFT are more complex than originally envisaged.

## **Retrograde IFT and Human Ciliopathies**

Defects in the HC that powers retrograde IFT (DYNC2H1) have been demonstrated to cause ciliopathies in humans, and have been particularly associated with short rib polydactyly and asphyxiating thoracic dystrophy (Jeune's syndrome) (Dagoneau et al., 2009; Merrill et al., 2009). Furthermore, HC mutations that result in premature termination occur only in the heterozygous state suggesting that the phenotype is hypomorphic (Schmidts et al., 2013). Mutations in the NEK1 kinase and the core IFT component IFT80 also result in the failure of ciliary assembly and lead to similar phenotypes (Beales et al., 2007; Thiel et al., 2011) although DYNC2H1 mutations are considered to be the major cause (Baujat et al., 2013). Our demonstration that lack of the WD60/FAP163 IFT dynein IC results in a severe ciliary assembly defect in planaria due to the failure of retrograde IFT thus raised the clear possibility that in vertebrates, mutations in this gene might cause phenotypes similar to those observed for DYNC2H1, NEK1 and IFT80. Indeed, a very recent report indicates that point mutations that result in truncation of WDR60 prior to the WD-repeat domain result in both short rib polydactyly and Jeune's syndrome (McInerney-Leo et al., 2013).

## **Materials and Methods**

### **Bioinformatics**

*Chlamydomonas* FAP163 was classified as a close relative of D1bIC/FAP133 (WD34 in vertebrates) by the PANTHER (*Protein Analysis THrough Evolutionary Relationships*) clustering algorithm (<http://www.pantherdb.org>). Orthologous sequences from other organisms closely related to *Chlamydomonas* FAP163 were identified using BLAST. Alignments with known dynein-associated WD-repeat proteins were generated using CLUSTALW (<http://www.clustal.org>), and the output used to construct a neighbor-joining phylogenetic tree.

The secondary structure of WD60 was assessed using PREDICTPROTEIN (<http://www.predictprotein.org>). Identification of WD-repeat domains, coiled coil regions and the repetitive N-terminal segment was performed using SMART (<http://smart.embl-heidelberg.de>), COILS ([http://embnet.vital-it.ch/software/COILS\\_form.html](http://embnet.vital-it.ch/software/COILS_form.html)) and DOTLET (<http://www.ikp.unibe.ch/dotlet/dotlet.html>), respectively. The IQ motif was identified using the Eukaryotic Linear Motif Resource (<http://elm.eu.org>).

### ***Chlamydomonas* Strains and Culture**

*Chlamydomonas reinhardtii* wildtype strains cc124/cc125 and mutants *fla10*, *oda1*, *oda2*, *oda3*, *oda4*, *oda5*, *oda6*, *ida1*, *ida4*, *pf14* and *pf18* were grown in R medium bubbled with 95% air/5% CO<sub>2</sub> on 15:9 hour light/dark cycle.

### **Preparation of $\alpha$ -FAP163 Antibody**

Sequence encoding the C-terminal 264 residues of FAP163 was obtained by PCR from first strand cDNA and subcloned into the pMAL-c2 vector across the XmnI/XbaI restriction sites resulting in a C-terminal fusion to maltose-binding protein (MBP). Expression of the MBP fusion protein in *Escherichia coli* BL21(DE3) was induced with 1 mM isopropylthiogalactoside and the protein subsequently purified by chromatography on an amylose affinity column. The purified fusion protein was then used as the immunogen for antibody production in rabbit CT295 (Covance Immunology Services, Denver, PA). Specific antibody was obtained from the resulting immune serum by blot affinity using the FAP163 C-terminal domain after removal of the MBP fusion partner.

Other antibodies used were rabbit polyclonals CT248, R4058 and R5932 against D1bIC(FAP133), LC8 and LC1, respectively (King and Patel-King, 1995; Benashski *et al.*,

1999; Rompolas *et al.*, 2007) and mouse monoclonal 1878A that reacts specifically with IC1 (King *et al.*, 1985).

### **Flagella Preparation and Biochemical Analysis of *Chlamydomonas* FAP163**

*Chlamydomonas* strains were deflagellated with dibucaine and flagella purified by standard methods (King, 1995). Flagella (~3 mg/ml) were demembrated with 1% Igepal CA-630 in 30 mM Hepes pH 7.5, 5 mM MgSO<sub>4</sub>, 0.5 mM EDTA, 25 mM KCl (HMEK buffer). Alternatively, matrix components were released by three rounds of freeze-thaw as described previously (Rompolas *et al.*, 2007). Chromatographic separations were performed using an ÄKTApurifier-10 workstation. Flagellar matrix samples were fractionated by gel filtration in a Superose 6 10/300 column equilibrated in HMEK, or were applied to a HR5/5 Mono Q ion exchange column equilibrated with 20 mM Tris.Cl pH 8.0 and then eluted using a 0-1 M NaCl gradient. For the *fla10* temperature-shift experiment, pre-warmed sucrose-containing buffers and dibucaine were added to the cell suspensions so that deflagellation occurred at the appropriate temperature. Subsequent flagella purification steps were performed at 4°C. Samples were electrophoresed in 10% or 12.5% SDS-polyacrylamide gels and either stained with Coomassie brilliant blue or transferred to nitrocellulose, stained with Reactive Brown 10 and subsequently probed with various antibodies (see above) followed by horseradish peroxidase-conjugated secondary antibodies. Chemiluminescence was detected using ECL reagent and a LAS4000 cooled CCD digital imaging system (both from GE systems, Piscataway, NJ).

### **RNA Interference in Planaria**

Colonies of the planarian *Schmidtea mediterranea* were maintained and expanded as detailed previously (Rompolas and King, 2009). RNA interference (RNAi) was performed essentially as described previously (Newmark *et al.*, 2003). In brief, to generate the RNAi

construct, a 325 bp region of the WD60 coding sequence was obtained from first-strand cDNA by PCR and then subcloned across the XbaI/XhoI sites in plasmid L4440 that contains opposing T7 promoters. Double-strand (ds)RNA production was induced following transformation into *E. coli* strain HT115(DE3) that lacks RNase III and thus cannot degrade dsRNA. Bacteria containing dsRNA or the empty vector control were fed to groups of planaria every 2-3 days for several weeks. The mRNA levels of the target and control genes were assessed by RT-PCR using standard methods.

### **Video Microscopy**

Videos of planarian movement in 9-cm petri dishes were taken with a CCD color camera (DFK 31BU03; The Imaging Source, Charlotte, NC) using IC-capture ver2.2 software at 30 fps and decompiled using Blaze Media Pro. Images illustrating the paths taken by individual animals were constructed by overlaying every 100<sup>th</sup> frame from the decompiled video using Adobe Photoshop CS4. The 20× real-time movie illustrating this motion was generated by decimating the video frames by twenty while keeping the frame rate at 30fps using VirtualDub. Individual movies were then cropped, combined and labeled using ImageJ. Higher magnification real-time videos were taken using a Canon Powershot SD880, converted with SUPER(C) and processed, cropped and combined with VirtualDub and ImageJ.

To visualize ciliary beating, individual planaria were confined on a microscope slide by parafilm spacers or held immobile using a Commodore Compressor (VU e-innovations, Nashville, TN). Cilia at the anterior dorsal-ventral junction were imaged using an Olympus BX51 microscope equipped with differential interference contrast optics and either a Plan Apo 60×/1.40 n.a. or 100×/1.35 n.a. oil objective lens. Videos of ciliary motility were captured using an X-PRI F1 digital camera (AOS Technologies, Baden Daetwil, Switzerland) at 250 fps. The

video frame rate, compression, cropping and brightness/contrast were adjusted using VirtualDub and ImageJ. Videos were decompiled with Blaze Media Pro.

### **Scanning and Transmission Electron Microscopy**

For scanning EM, planaria were treated overnight with relaxant solution (1% HNO<sub>3</sub>, 0.85% formaldehyde, 50 mM MgSO<sub>4</sub>), rinsed with buffer and then fixed with 2.5% glutaraldehyde in 50 mM Na cacodylate pH 7.4. Subsequently, samples were post-fixed with 1% OsO<sub>4</sub> and dehydrated through an ethanol series. Following critical point drying, planaria were mounted, sputter coated with gold and imaged in a JEOL JSM5900 scanning EM.

To visualize internal ciliary structure, the anterior region of individual planarians was excised, and fixed with 1% glutaraldehyde in PBS for 15 mins and then with 1% glutaraldehyde in 50 mM Na cacodylate for 1 h; the fixative solution was changed every 10-15 mins to avoid it becoming exhausted due to the large quantity of mucus on the planarian ventral surface. Samples were post-fixed with 1% OsO<sub>4</sub>, stained *en bloc* with 1% methanolic uranyl acetate, dehydrated through ethanol and embedded in Epon. Thin sections were post-stained with uranyl acetate and lead citrate, and imaged using a Hitachi H-7650 transmission EM operating at 80 kV.

### **Acknowledgements**

We thank Dr Panteleimon Rompolas (Yale University School of Medicine) for many helpful discussions and Maya Yankova (UCHC) for her assistance with electron microscopy. We are also grateful to Dr Carol Wicking (University of Queensland) for providing a preprint describing human WDR60 mutations prior to publication. This study was supported by grant GM051293 from the National Institutes of Health (to S.M.K). E.F.Y.H. was supported in part by the Jane Coffin Childs Memorial Research Fund and the NIGMS Center for Systems Biology (GM068763).

## Footnotes

1. Abbreviations used: EM, electron microscopy; FAP, flagellar-associated protein; HC, heavy chain; IC, intermediate chain; IFT, intraflagellar transport; LC, light chain; LIC, light intermediate chain; MBP, maltose-binding protein.
2. As we demonstrate that FAP163 is a dynein IC involved in retrograde IFT, we propose that the *Chlamydomonas* gene name be reassigned as *DIC6* in accordance with the unified nomenclature for ciliary dyneins (Hom et al., 2011). Previously, the FAP133 protein was assigned as D1bIC (Hom et al., 2011). As it is now clear that this dynein contains two different ICs, we suggest that FAP163 and FAP133 be renamed as D1bIC1 and D1bIC2, respectively.

## References

- Baujat, G.v., Huber, C.I., El Hokayem, J., Caumes, R., Do Ngoc Thanh, C., David, A., Delezoide, A.-L., Dieux-Coeslier, A., Estournet, B., Francannet, C., Kayirangwa, H., Lacaille, F., Le Bourgeois, M., Martinovic, J., Salomon, R.m., Sigaudy, S., Malan, V.r., Munnich, A., Le Merrer, M., Le Quan Sang, K.H., and Cormier-Daire, V.R. (2013). Asphyxiating thoracic dysplasia: clinical and molecular review of 39 families. *J Medical Genetics* 50 91-98
- Beales, P.L., Bland, E., Tobin, J.L., Bacchelli, C., Tuysuz, B., Hill, J., Rix, S., Pearson, C.G., Kai, M., Hartley, J., Johnson, C., Irving, M., Elcioglu, N., Winey, M., Tada, M., and Scambler, P.J. (2007). IFT80, which encodes a conserved intraflagellar transport protein, is mutated in Jeune asphyxiating thoracic dystrophy. *Nature Genetics* 39, 727-729.

Benashski, S.E., Patel-King, R.S., and King, S.M. (1999). Light chain 1 from the *Chlamydomonas* outer dynein arm is a leucine-rich repeat protein associated with the motor domain of the  $\gamma$  heavy chain. *Biochemistry* 38, 7253-7264.

Cole, D. (2009). Intraflagellar transport. In: The *Chlamydomonas* Source Book, vol. 3: Cell Motility and Behavior, ed. G.B. Witman, San Diego: Elsevier, 71-113.

Cole, D.G. (2003). The intraflagellar transport machinery of *Chlamydomonas reinhardtii*. *Traffic* 4, 435-442.

Dagoneau, N., Goulet, M., Geneviève, D., Sznajder, Y., Martinovic, J., Smithson, S., Huber, C.I., Baujat, G.v., Flori, E., Tecco, L., Cavalcanti, D., Delezoide, A.-L., Serre, V.r., Le Merrer, M., Munnich, A., and Cormier-Daire, V.r. (2009). DYNC2H1 Mutations cause asphyxiating thoracic dystrophy and short rib-polydactyly syndrome, type III. *Amer J Human Genetics* 84, 706-711.

Efimenko, E., Bubb, K., Mak, H.Y., Holzman, T., Leroux, M.R., Ruvkun, G., Thomas, J.H., and Swoboda, P. (2005). Analysis of *xbx* genes in *C. elegans*. *Development* 132, 1923-1934.

Hao, L., Efimenko, E., Swoboda, P., and Scholey, J. (2011). The retrograde IFT machinery of *C. elegans* cilia: two IFT dynein complexes? *PLoS One*, e20995.

Hildebrandt, F., Benzing, T., and Katsanis, N. (2011). Ciliopathies. *New England Journal of Medicine* 364, 1533-1543.

Hoh, R., Stowe, TR, Turk, E, and Stearns, S. (2012). Transcriptional program of ciliated epithelial cells reveals new cilium and centrosome components and links to human disease. *PLoS One* 7, e52166.

Hom, E., Witman, G.B., Harris, E.H., Dutcher, S.K., Kamiya, R., Mitchell, D.R., Pazour, G.J., Porter, M.E., Sale, W.S., Wirschell, M., Yagi, T., and King, S.M. (2011). A unified taxonomy for ciliary dyneins. *Cytoskeleton* 68, 555-565.

King, S.M. (1995). Large-scale isolation of *Chlamydomonas* flagella. *Methods Cell Biol* 47, 9-12.

King, S.M. (2012). Composition and assembly of axonemal dyneins. In: *Dyneins: Structure, Biology and Disease*, ed. S.M. King, Waltham, MA: Elsevier, 209-243.

King, S.M., Otter, T., and Witman, G.B. (1985). Characterization of monoclonal antibodies against *Chlamydomonas* flagellar dyneins by high-resolution protein blotting. *Proc Natl Acad Sci USA* 82, 4717-4721.

King, S.M., and Patel-King, R.S. (1995). The  $M_{(r)} = 8,000$  and 11,000 outer arm dynein light chains from *Chlamydomonas* flagella have cytoplasmic homologues. *J Biol Chem* 270, 11445-11452.

King, S.M., Wilkerson, C.G., and Witman, G.B. (1991). The  $M_r$  78,000 intermediate chain of *Chlamydomonas* outer arm dynein interacts with alpha-tubulin *in situ*. *J Biol Chem* 266, 8401-8407.

Li, J.B., Gerdes, J.M., Haycraft, C.J., Fan, Y., Teslovich, T.M., May-Simera, H., Li, H., Blacque, O.E., Li, L., Leitch, C.C., Lewis, R.A., Green, J.S., Parfrey, P.S., Leroux, M.R., Davidson, W.S., Beales, P.L., Guay-Woodford, L.M., Yoder, B.K., Stormo, G.D., Katsanis, N., and Dutcher, S.K. (2004). Comparative genomics identifies a flagellar and basal body proteome that includes the BBS5 human disease gene. *Cell* 117, 541-552.

Lo, K.W.-H., Kan, H.-M., and Pfister, K.K. (2006). Identification of a novel region of the cytoplasmic dynein intermediate chain important for dimerization in the absence of the light chains. *J. Biol. Chem.* 281, 9552-9559.

McInerney-Leo, A.M., Schmidts, M., Cortes, C.R., Leo, P.J., Gner, B., Courtney, A.D., Gardiner, B., Harris, J.A., Lu, Y., Marshall, M., UK10K Consortium, Scambler, P.J., Beales,

P.P., Brown, M.A., Zankl, A., Mitchison, H., Duncan, E.L., and Wicking, C. (2013). Exome sequencing identifies WDR60 as a novel causative gene in short rib polydactyly and Jeune syndromes. *Am. J. Hum. Genet.* in press.

Merrill, A.E., Merriman, B., Farrington-Rock, C., Camacho, N., Sebald, E.T., Funari, V.A., Schibler, M.J., Firestein, M.H., Cohn, Z.A., Priore, M.A., Thompson, A.K., Rimoin, D.L., Nelson, S.F., Cohn, D.H., and Krakow, D. (2009). Ciliary abnormalities due to defects in the retrograde transport protein DYNC2H1 in short-rib polydactyly syndrome. *Amer J Human Genetics* 84, 542-549.

Mitchell, D.R., and Kang, Y. (1991). Identification of *oda6* as a *Chlamydomonas* dynein mutant by rescue with the wild-type gene. *J Cell Biol* 113, 835-842.

Mitchell, D.R., and Rosenbaum, J.L. (1986). Protein-protein interactions in the 18S ATPase of *Chlamydomonas* outer dynein arms. *Cell Motil Cytoskeleton* 6, 510-520.

Newmark, P., Reddien, P., Cebria, F., and Sanchez-Alvarado, A. (2003). Ingestion of bacterially expressed double-stranded RNA inhibits gene expression in planaria. *Proc Natl Acad Sci USA* 100, 11861-11865.

Nyarko, A., Hare, M., Hays, T.S., and Barbar, E. (2004). The intermediate chain of cytoplasmic dynein is partially disordered and gains structure upon binding to light-chain LC8 *Biochemistry* 43, 15595-15603.

Pan, X., Ou, G., Civelekoglu-Scholey, G., Blacque, O.E., Endres, N.F., Tao, L., Mogilner, A., Leroux, M.R., Vale, R.D., and Scholey, J.M. (2006). Mechanism of transport of IFT particles in *C. elegans* cilia by the concerted action of kinesin-II and OSM-3 motors. *J Cell Biol* 174 1035-1045

Pazour, G., Agrin, N., Leszyk, J., and Witman, G. (2005). Proteomic analysis of a eukaryotic flagellum. *J Cell Biol* *170*, 103-113.

Pazour, G.J., Dickert, B.L., Vucica, Y., Seeley, E.S., Rosenbaum, J.L., Witman, G.B., and Cole, D.G. (2000). *Chlamydomonas* IFT88 and its mouse homologue, polycystic kidney disease gene *tg737*, are required for assembly of cilia and flagella. *J Cell Biol* *151*, 709-718.

Pazour, G.J., Dickert, B.L., and Witman, G.B. (1999). The DHC1b (DHC2) isoform of cytoplasmic dynein is required for flagellar assembly. *J Cell Biol* *144*, 473-481.

Pazour, G.J., Wilkerson, C.G., and Witman, G.B. (1998). A dynein light chain is essential for the retrograde particle movement of intraflagellar transport (IFT). *J Cell Biol* *141*, 979-992.

Perrone, C., Tritschler, D., Taulman, P., Bower, R., Yoder, B., and Porter, M. (2003). A novel dynein light intermediate chain colocalizes with the retrograde motor for intraflagellar transport at sites of axoneme assembly in *Chlamydomonas* and mammalian cells. *Mol Biol Cell* *14*, 2041-2056.

Porter, M.E., Bower, R., Knott, J.A., Byrd, P., and Dentler, W. (1999). Cytoplasmic dynein heavy chain 1b is required for flagellar assembly in *Chlamydomonas*. *Mol Biol Cell* *10*, 693-712.

Rompolas, P., and King, S.M. (2009). *Schmidtea mediterranea*: a model system for analysis of motile cilia. *Methods Cell Biol* *93*, 81-98.

Rompolas, P., Patel-King, R.S., and King, S.M. (2010). An outer arm dynein conformational switch is required for metachronal synchrony of motile cilia in planaria. *Mol Biol Cell* *21*, 3669-3679.

Rompolas, P., Pedersen, L., Patel-King, R.S., and King, S.M. (2007). *Chlamydomonas* FAP133 is a dynein intermediate chain associated with the retrograde intraflagellar transport motor. *J Cell Sci* *120*, 3653-3665.

Rosenbaum, J.L., and Witman, G.B. (2002). Intraflagellar transport. *Nature Reviews Molecular and Cellular Biology* 3, 813-825.

Schmidts, M., Arts, H.H., Bongers, E.M.H.F., Yap, Z., Oud, M.M., Antony, D., Duijkers, L., Emes, R.D., Stalker, J., Yntema, J.-B.L., Plagnol, V., Hoischen, A., Gilissen, C., Forsythe, E., Lausch, E., Veltman, J.A., Roeleveld, N., Superti-Furga, A., Kutkowska-Kazmierczak, A., Kamsteeg, E.-J., ElÄşioÄŸlu, N., van Maarle, M.C., Graul-Neumann, L.M., Devriendt, K., Smithson, S.F., Wellesley, D., Verbeek, N.E., Hennekam, R.C.M., Kayserili, H., Scambler, P.J., Beales, P.L., Knoers, N.V., Roepman, R., and Mitchison, H.M. (2013). Exome sequencing identifies DYNC2H1 mutations as a common cause of asphyxiating thoracic dystrophy (Jeune syndrome) without major polydactyly, renal or retinal involvement. *J Medical Genetics* 50 309-323

Signor, D., Wedaman, K.P., Orozco, J.T., Dwyer, N.D., Bargmann, C.I., Rose, L.S., and Scholey, J.M. (1999). Role of a class DHC1b dynein in retrograde transport of IFT motors and IFT raft particles along cilia, but not dendrites, in chemosensory neurons of living *Caenorhabditis elegans*. *J Cell Biol* 147, 519-530.

Snow, J.J., Ou, G., Gunnarson, A.L., Walker, M.R.S., Zhou, H.M., Brust-Mascher, I., and Scholey, J.M. (2004). Two anterograde intraflagellar transport motors cooperate to build sensory cilia on *C. elegans* neurons. *Nature Cell Biol* 6, 1109-1113.

Tang, W.J., Bell, C.W., Sale, W.S., and Gibbons, I.R. (1982). Structure of the dynein-1 outer arm in sea urchin sperm flagella. I. Analysis by separation of subunits. *J Biol Chem* 257, 508-515.


Thiel, C., Kessler, K., Giessel, A., Dimmler, A., Shalev, S.A., von der Haar, S., Zenker, M., Zahnleiter, D., Stöss, H., Beinder, E., Abou Jamra, R., Ekici, A.B., Schröder-Kress, N., Aigner,

T., Kirchner, T., Reis, A., Brandstätter, J.H., and Rauch, A. (2011). NEK1 mutations cause short-rib polydactyly syndrome type Majewski. *Amer J Human Genetics* 88, 106-114.

Wickstead, B., and Gull, K. (2006). A holistic kinesin phylogeny reveals new kinesin families and predicts protein functions. *Mol Biol Cell* 17, 1734-1743.

Wickstead, B., and Gull, K. (2012). Evolutionary biology of dyneins. In: *Dyneins: Structure, Biology and Disease*, ed. S.M. King, Waltham, MA: Elsevier, Inc, 89-121.

Witman, G.B. (2012). Dynein and intraflagellar transport. In: *Dyneins: Structure, Biology and Disease*, ed. S.M. King, Boston: Elsevier, 395-421.



**Fig. 1 Phylogeny and Domain Organization of WD60/FAP163**


a) The sequences of dynein-associated WD-repeat components, as well as mammalian, planarian and algal WD60/FAP163 proteins were aligned using CLUSTALW, and a neighbor-joining tree generated. All WD60/FAP163 sequences cluster together, and are most closely related to the *Chlamydomonas* retrograde dynein IC D1bIC(FAP133) and its human ortholog (WD34). Sequences used were: *Homo sapiens* (Hs) WD60 (NP\_060521), *Rattus norvegicus* (Rn) WD60 (NP\_001178702), *Bos taurus* (Bt) WD60 (XP\_001788990), *S. mediterranea* (Sm) WD60 (mk4.000489.09.01), *C. reinhardtii* (Cr) FAP163 (XP\_001702532), *Volvox carteri* (Vc) FAP163 (XP\_002954132), *H. sapiens* WD34 (AAH01614), *C. reinhardtii* D1bIC/FAP133 (BC133909), *C. reinhardtii* IC2 (P27766), *Strongylocentrotus purpuratus* (Sp) IC3 (XP\_796667), *C. reinhardtii* IC140 (EDP01123), *C. reinhardtii* Lis1 (ABG33844), *Mus*

*musculus* Lis1 (NP\_038653), *C. reinhardtii* IC1 (Q39578) and *St. purpuratus* IC2 (XP\_003724308).

b) The propensity for coiled coil formation within human WD60 was analyzed using the COILS algorithm. Two segments have a very high prediction for coiled coil formation.

c) Comparison of human WD60 against itself using DOTLET (matrix = BLOSUM62; window = 15) revealed that the N-terminal region is repetitive consisting of multiple charged segments with high similarity.

d) Domain organization of human WD60. The N-terminal region contains two coiled-coil segments embedded within a region predicted to be almost entirely  $\alpha$ -helical. The C-terminal domain consists of a series of predicted  $\beta$  strands; three canonical WD repeats were identified within this region using SMART (residues 686-725, 901-938 and 941-983). WD repeats are often highly degenerate and difficult to identify, however, visual inspection of the predicted secondary structure suggests there may be two additional poorly conserved repeats between the first and second canonical WD-repeats and a third prior to the first canonical repeat. An IQ motif is located between residues 432 and 450.




**Fig. 2 FAP163 is a Component of the Flagellar Matrix**

a) Flagellar proteins from the *oda1* mutant were separated in a 10% polyacrylamide gel and either stained with Coomassie blue (CBB) or blotted to nitrocellulose and probed with the CT295 antibody raised against FAP163. The antibody detects a single major band of M<sub>r</sub>89,000 and three very minor bands.

b) Flagella from wildtype *Chlamydomonas* (*cc125*) and mutants lacking outer arms (*oda1-oda6*), inner arms (*ida1* and *ida4*), radial spokes (*pf14*) or the central pair microtubule complex (*pf18*) were electrophoresed in 10% polyacrylamide gels and either stained with Coomassie blue

(*upper panels*) or blotted and probed with CT295 to detect FAP163 (*lower panels*). This protein was present in all the mutant flagella tested.

c) To determine whether FAP163 is associated with the axoneme or part of the flagella matrix, wildtype flagella were extracted with detergent (1% Igepal CA-630) to yield a supernatant containing the membrane and matrix components and a pellet containing the microtubular axoneme. Flagella were also fractionated by three rounds of freeze/thaw which releases the matrix components but leaves the membrane associated with the axoneme. The *upper panels* show the Coomassie blue-stained 10% polyacrylamide gels and the *lower panels* blots probed with CT295. This revealed that FAP163 is released by both detergent and freeze/thaw indicating that it is a component of the flagella matrix.




Patel-King et al.  
Figure 3

**Fig. 3 FAP163 and D1b1C(FAP133) Associate with Each Other and**


### **the LC8 Dynein Light Chain**

- a) Chromatogram of an *odal* freeze/thaw extract fractionated by gel filtration on a Superose 6 10/300 column. The bar indicates the region analyzed by SDS-PAGE. The *lower panels* show these fractions separated in 10% polyacrylamide gels and stained with Coomassie blue, or blotted and probed with CT295, CT248 and R4058 to reveal FAP163, D1bIC(FAP133) and LC8, respectively; samples for analysis of LC8 content were separated in 12.5% gels.
- b) Chromatogram of a wildtype (cc125) freeze/thaw extract that was applied to a Mono Q anion exchange column and eluted with a 0-1 M gradient of NaCl. The *upper panel* shows the resulting chromatogram and the *lower panels* the immunoblots corresponding to the region on the chromatogram indicated by the bar. D1bIC(FAP133) and FAP163 co-elute in two separate peaks.



**Fig. 4 FAP163 is Trafficked by the Intraflagellar Transport Machinery**

Wildtype (*cc125*) and *fla10* cells were maintained for 3 hours at either the permissive (22°C) or restrictive (32°C) temperature, flagella isolated and then electrophoresed in 10% or 12.5% polyacrylamide gels. The *upper panel* shows the Coomassie blue-stained 10% gel and the *lower panels* immunoblots probed with antibodies CT248, CT295, R5932 and 1878A to detect D1bIC(FAP133), FAP163, LC1 and IC1, respectively. Both D1bIC and FAP163 levels were reduced in *fla10* flagella at the restrictive temperature suggesting that they are actively trafficked by the IFT machinery whereas LC1 and IC1 were not.



**Fig. 5** *Smed-wd60*(RNAi) Planaria Exhibit Reduced Gliding Velocity and an Altered Mode of Locomotion

a) RT-PCR analysis of mRNA levels in untreated control planaria and in animals fed either with the empty L4440 vector or dsRNA from the *Smed-wd60*(RNAi) construct are shown; the first-strand cDNA was prepared using a random hexamer as primer and the reaction was

continued for 25 cycles. Planaria fed with the WD60 dsRNA have unchanged levels of mRNAs encoding actin, IFT88, and outer arm dynein IC2. However, two RT-PCR products (325 and 502 bp) were detected for WD60. Sequence analysis revealed that both encoded the same region of WD60 but that the larger one also contained two unspliced introns. Levels of the 325 bp product were greatly reduced in planaria fed with WD60 dsRNA whereas levels of the 502 bp product were unchanged. The 502 bp product was not obtained following treatment of the RNA sample with DNaseI prior to PCR (not shown), indicating that it derives from genomic DNA.


b) Map of the two RT-PCR products for WD60 shown in (a) indicating the location of the two introns.

c) Products obtained by RT-PCR after 40 cycles following DNaseI-treatment of the mRNA sample and first-strand cDNA synthesis using an oligo-dT primer. The two additional products derive from partially spliced polyadenylated mRNA for WD60.

d) Bar chart illustrating the velocity of movement of control and *Smed-wd60(RNAi)* planaria. The controls moved by cilia-driven gliding at a rate of  $1.48 \pm 0.21$  mm/s (mean  $\pm$  std. dev.;  $n = 7$ ), whereas the knockdown animals utilize waves of muscle contraction and traveled at  $0.55 \pm 0.19$  mm/s ( $n = 13$ ).


e) The tracks taken by individual planaria over a 30 sec period are shown; these images were prepared from the decompiled videos by overlaying every 100<sup>th</sup> frame using Adobe Photoshop (and see supplemental movie S1).

f) Mode of planarian movement. Controls move with a smooth cilia-driven gliding motion whereas *Smed-wd60(RNAi)* animals employ peristaltic waves of muscle contraction. These images were taken from the decompiled combined video (see supplemental movie S2).




**Fig. 6** *Smed-wd60(RNAi)* Planaria have Defects in Ciliary Assembly

Scanning electron micrographs of the ventral surface of control (L4440; *panels a and c*) and *Smed-wd60(RNAi)* (*panels b, d and e*) animals are shown. The control planaria have many closely apposed cilia that are ~10 μm in length. In contrast, the ventral surface of *Smed-wd60(RNAi)* animals have large areas that are almost devoid of full-length cilia, but rather have many ciliary stubs that are <0.5 μm long. The bars are 100 μm (*panels a and b*), 10 μm (*panels c and d*) and 2 μm (*panel e*).



**Fig. 7 Cilia and Ciliary Stubs Present on *Smed-wd60(RNAi)* Planaria Contain Large Quantities of IFT Particles**

Transmission electron micrographs of cilia on the ventral surface of control animals (L4440; *left panels*), and cilia and ciliary stubs of *Smed-wd60(RNAi)* planaria (*all other panels*). The accumulation of electron-dense material (presumably IFT particles) between the doublet microtubules and ciliary membrane is clearly evident in both longitudinal and transverse sections of the knock-down animals; in contrast, this accumulated material is absent from control cilia. The bar is 100 nm and applies to all panels.



**Fig. 8 Cilia Assembled by *Smed-wd60(RNAi)* Planaria have an Aberrant Beat**

a) Sequential frames from the decompiled videos taken at 250 fps of ciliary beating for control and *Smed-wd60(RNAi)* planaria. The numbers at *upper left* in each frame indicate the time in msec. Cilia in the knockdown animals are capable of motility although their beat is uncoordinated and chaotic. In some cases, the cilia twitch from the base and do not show evidence of effective power and effective strokes (and see supplemental movies S3 and S4).

b) Kymographs (0.5 sec) illustrating the beat frequency of control and *Smed-wd60(RNAi)* planaria. The arrows on the two panels indicate each beat cycle.

# Supplemental Materials

*Molecular Biology of the Cell*

Patel-King et al.



